

Increased Ammonia leads to...

- ❑ Less profits
- ❑ Poor FCR
- ❑ Decreased weight gain
- ❑ Respiratory tract damage
- ❑ Eye damage
- ❑ High mortality
- ❑ Increased Ascites
- ❑ Increased Disease Severity
- ❑ Increased vaccination Failures/reactions

Liiteron

The Litter Conditioner

Recommended Usage :

Birds on deep litter :

- ❑ Control ammonia – spread 5 Kg – 10 Kg Liiteron per 1000 Sq. Ft. of floor space.
- ❑ Break cycle of reinfection for E.Coli, Salmonella, Pseudomonas, Campylobacter – spread 10 Kg Liiteron per 1000 Sq. Ft. floor space. Repeat every 30 days.

Birds in Cages :

Spread Liiteron 20 Kg per 1000 Sq. Ft. on the floor/litter. Liiteron usage may be repeated every 30 days or earlier depending upon litter quality.

Repeat use of litter :

After birds have been taken out, remove the litter and disinfect(Maxii Guard) the shed. Remove all cake formation from litter. Mix 10 Kg Liiteron in litter to be spread over every 1000 Sq. Ft. area. After similar action the litter can be used again and again upto 10 times.


Presentation : Packs of 10 Kg & 25 Kg

Manufactured by :

 **Guybro Chemical**

201, Maruti Chamber, Fun Republic Lane,
Veera Desai Road, Andheri(W), Mumbai 400 53.
Tel. No. : 91-22-40546800 Fax No. : 91-22-40546811
Email : guybrochemical@hotmail.com
Website : www.toxinbinder.com

Control Poultry House Ammonia Litter Bacteria And Improve Profits



Liiteron
The Litter Conditioner