



SPI LAC

HARNESSING THE POWER OF
AQUA FEEDS

With
SPIRULINA*

FEED SPILAC FOR

- Building up Resistance
- Faster growth
- Increasing Biomass
- Lowering FCR & Increasing Profits

* "The greatest food available on Earth"
(W.H.O.)

FEED SPILAC FROM DAY ONE TO MARKETING

SPIILAC[®]

With

SPIRULINA

"The greatest food available on Earth"

World Health Organisation (W.H.O.)

COMPOSITION :

Each Kg Contains :		Mineral Content :			
Lactobacillus sporogenes	33,200 Million cfu	Calcium	50 mg	Methionine	70 mg
Liver extract	5320 mg	Iron	7.5 mg	Phenylalanine	140 mg
Yeast extract (Saccharomyces cerevisiae)	1250 mg	Zinc	0.15 mg	Threonine	160 mg
Alpha amylase	5000 mg	Phosphorus	45 mg	Tryptophan	45 mg
Ascorbic acid coated	500 mg	Magnesium	40 mg	Valine	200 mg
Protein	100 gm	Copper	0.06 mg	Alanine	235 mg
SPIRULINA	5320 mg	Sodium	30 mg	Arginine	215 mg
With following ingredient		Potassium	80 mg	Glutamic acid	455 mg
Vitamin Content :		Manganese	0.25 mg	Histidine	50 mg
Vit E (α - tocopherol)	0.51.U.	Chromium	0.014 mg	Proline	135 mg
Vit A (β - carotene)	15003 I.U.	Germanium	0.003 mg	Serine	160 mg
Vit B ₁ (thiamine)	0.155 mg	Selenium	0.001 mg	Tyrosine	150 mg
Vit B ₂ (riboflavin)	0.175 mg	Natural Pigments :		Fatty Acid	
Vit B ₃ (niacin)	0.73 mg	Phycocyanin(blue)	750-1000 mg	C14.0 Myristic	0.5 mg
Vit B ₆ (pyridoxine)	0.04 mg	Chlorophyll(green)	5.75 mg	C16.0 Palmitic	122 mg
Vitamin B ₁₂	0.016 mg	Carotenoids(orange)	18.5 mg	C16.1 Palmitoleic	16.5 mg
Folacin	0.005 mg	Beta carotene	7 mg	C17.0 Heptadecanoic	1 mg
Inositol	3.24 mg	Amino acids :		C18.0 Stearic	4 mg
		Isoleucine	175 mg	C18.1 Oleic	6 mg
		Leucine	270 mg	C18.2 Linolenic	48.5 mg
		Lysine	145 mg	C18.3 Gamma-Linoleic	67.5 mg
				C Others	7 mg
				Feed attractant and base	Q.S.

INDICATIONS :

- GROWTH PHASE :** 2-4 weeks.
- STRESS CONDITIONS :**
 - Sudden changes in weather
 - Extreme heat or cold.
 - Change in feed.
 - Disease outbreak in the vicinity
- FINISHING STAGES :** 2-4 weeks.
- Helps overcome stress due to intensive and Semi-Intensive system of Scampiculture / Aquaculture practices.

FEEDING RECOMMENDATION :

- Fish : 1Kg per Tonne of feed
 Shrimps/Prawns : 2 Kg per Tonne of Mash feed.
 5 Kg per Tonne of Pellet feed.
 (5 g per Kg)
- For best result, continuous use is recommended.

Presentation : Available in 1 Kg. Polyjar & 5 Kg. Bucket

SPIILAC[®]

A COMPLETE NUTRITION

ENRICHED WITH SPIRULINA AND PREDIGESTED PROTEIN
 ENSURES PRODUCTIVITY AND PROFITS.



Guybro Chemical

Admn. Off. : 201, Maruti Chambers, Off. Veera Desai Road, Fun Republic Lane, Andheri (W), Mumbai - 400 053.

Tel. : +91-22-4054 6828 • Contact info : guybrochemical@hotmail.com

[®] Registered Trademark of Guybro Chemical